## LASIK Post Surgery Drop Schedule

Day of Surgery	Once Home	After Nap	Dinner	Bedtime
Zymaxid (Gatifloxacin)	Х	×	×	×
Pred Forte (Prednisolone) (shake well)	Х	Х	Every 2 hrs. while awake	
Preservative Free Artificial Tears	Х	×	Every 30 mins. while awake	

Instill drops 5 minutes apart (1 drop each eye) as indicated by the X. Use 2 drops in each eye of Artificial Tears.

Days 1 - 2	Breakfast	Lunch	Dinner	Bedtime	
Zymaxid (Gatifloxacin)	Х	Х	Х	Х	
Pred Forte (Prednisolone) (shake well)	Every 2 hrs. while awake				
Preservative Free Artificial Tears	Every 30 mins. while awake				

Instill drops 5 minutes apart (1 drop each eye) as indicated by the X. Use 2 drops in each eye of Artificial Tears.

Days 3 - 7	Breakfast	Lunch	Dinner	Bedtime
Zymaxid (Gatifloxacin)	Х	Х	Х	Х
Pred Forte (Prednisolone) (shake well)	Х	Х	Х	Х
Preservative Free Artificial Tears	Every 30 mins. while awake			

Continue Preservative Free Artificial Tears (vials) for 30 days after surgery. Use 2 drops in each eye (use vial until empty) then throw away.

Please call (843) 881-3937 if you have any questions or concerns.